arXiv:1107.4964v1 [quant-ph] 25 Jul 2011

Coherently manipulating cold ions in separated traps by their vibrational couplings II: non-adiabatic case

Miao Zhang[1] and L. F. Wei*[1,2,3]
[1]Quantum Optoelectronics Laboratory, School of Physics and Technology,
Southwest Jiaotong University, Chengdu 610031, China
[2]State Key Laboratory of Optoelectronic Materials and Technologies,
School of Physics and Engineering, Sun Yat-sen University, Guangzhou 510275, China
[3]State Key Laboratory of Functional Materials for Informatics,
Shanghai Institute of Microsystem and Information Technology,
Chinese Academy of Sciences, Shanghai 200050 China

(Dated: July 26, 2011)

In the previous paper [M. Zhang and L. F. Wei, Phys. Rev. A **83**, 064301 (2011)] we proposed an approach to implement the universal quantum gates between two ions, confined individually in the separated traps, by adiabatically manipulating the ionic vibrations. Here, we relax the previous adiabatic requirements and show that the desirable couplings between the ions in different traps can be still implemented by simply using a non-adiabatic laser pulse. The potential wells of the ions are unchanged, but the ionic vibrations are driven for resonance or large-detuning by the applied laser beams. Consequently, the desirable quantum gates between the ions could be implemented conveniently by using only a few of laser pulses. The proposal could also be generalized to the fast quantum computation with the scalable ion-traps.

PACS number(s): 03.67.Lx, 42.50.Dv, 37.10.Ty

Among the proposed physical systems for building the future quantum computer [1–5], the system of trapped ions [6] is currently one of the most hopeful candidates. Indeed, the entanglement up to fourteen trapped ions has been experimentally realized [7]. In the usual implementations a string of ions are trapped in a single trap, and their center-of-mass (CM) mode acts as the data bus to couple the qubits encoded by the long-lived electronic levels [8, 9]. However, it should be a challenge to trap more ions in a single well and then coherently manipulate them precisely, as it is difficult to obtain the reasonable strong-radial-confinements to form a linear ion-string [10]. Furthermore, the speed of the laser-induced sideband excitation should be significantly reduced with the larger mass of an ion-chain.

Alternatively, the scheme of the segmented ion-traps could be an more practical way to implement the large-scale quantum computing network [11–13]. This is because that the ions can be relatively-easily confined in the separated potential wells, and their couplings can be realized by transporting the ions to any selected interaction region. Alternatively, if the separate wells are sufficiently close, then the trapped ions can be coupled directly via their Coulomb interaction, and the transportation of ions are avoided. Interestingly, such an idea has been demonstrated in recent experiments [14, 15], wherein two ions are confined in two potential wells separated by $40\,\mu$m [14] (or $54\,\mu$m [15]) and the ion-ion coupling is achieved up to $g \approx 10$ KHz (or $g \approx 7$ KHz).

In our recent paper [16] we have proposed an approach to generate the universal quantum gates between the two separately trapped ions. That idea is based on the experiments [14, 15]: the coupling between the ions was manipulated by controlling the potential wells (i.e., the voltages on the DC electrodes) to adiabatically tune ions' external vibrations into or out of resonance. Obviously, the adiabatic conditions limit the speed of the implemented quantum gates. Also, the exactly-controlling of the potential wells, by adjusting the voltages applied on the DC electrodes, should be very complex for the large-scale ion-traps.

In order to overcome these difficulties, in this brief report we propose an improved approach to implement the universal quantum gates [17] between the separately trapped ions. Instead of the previous adiabatic operations applied directly on the potential wells [14–16], here we use the laser beams to drive one of the trapped ions for controlling its interaction with another ion, let them be resonance or large-detuning. With the present switchable couplings the desirable coherent operations, e.g., the quantum gates, between the separately trapped ions could be performed conveniently by using only a few laser pulses. Hopefully, the present approach is more robust for the large-scale ion-traps.

We consider the same experimental configuration (shown in Fig. 1) as that in the previous works [14–16], i.e., two ions trapped individually in two potential wells separated by a distance d. The ion trap provides a pseudopotential such that the ions' oscillating frequencies along the axial direction (i.e., z axis in the Fig. 1) is much smaller than those along the radial directions (i.e., the x, y axis). As a consequence, only the quantized vibrational motion along the axial direction is considered, and the Hamiltonian describing the oscillating ions reads [14–16]

$$\hat{H}_{\rm ex} = \sum_{j=1}^{2} \hbar\nu_j(\hat{a}_j^{\dagger}\hat{a}_j + \frac{1}{2}) + \hat{V}_{\rm ii} \tag{1}$$

---

*weilianfu@gmail.com



FIG. 1: (Color online) Sketch for implementing switchable coupling between two ions trapped individually in two potential wells separated by a distance d. The ions are trapped initially in the large-detuning regime, i.e., they are decoupled at the beginning, then a laser pulse (applied to the ion 2) makes $\Delta = 0$ for generating a resonant coupling.

with

$$\hat{V}_{\rm ii} = \mathcal{K}\Bigg[\textstyle\sum_{j=1}^{2} \frac{(-1)^j \xi_j}{d}\hat{a}_j + \frac{\xi_j^2}{d^2}(\hat{a}_j^2 + \hat{a}_j\hat{a}_j^\dagger) \\ -\frac{2\xi_1\xi_2}{d^2}(\hat{a}_1\hat{a}_2 + \hat{a}_1\hat{a}_2^\dagger)\Bigg] + \text{H.c.} \tag{2}$$

being the ion-ion Coulomb interaction. Where, $\hat{a}_j^\dagger$ and $\hat{a}_j$ are the bosonic creation and annihilation operators of the vibrations of frequency ν_j. The parameters $\mathcal{K}$ and ξ_j are defined as $\mathcal{K} = q_1q_2/(4\pi\epsilon_0 d)$ and $\xi_j = \sqrt{\hbar/(2M_j\nu_j)}$, where M_j and q_j are the mass and charge of the ions, respectively.

For realizing a switchable coupling between the two trapped ions, without changing the eigenfrequencies v_j of the ions, we apply the laser beams to one of the ions, e.g., the ion 2. The Hamiltonian describing the laser-driving ion takes the well-known form [18, 19]

$$\hat{H}_{\rm L} = \frac{\hbar\omega_a}{2}\hat{\sigma}_z + \hbar\Omega_0(\hat{\sigma}_+ + \hat{\sigma}_-)\left[e^{i\eta(\hat{a}_2+\hat{a}_2^\dagger)-i\omega_l t - i\phi_l} + \text{H.c.}\right]. \tag{3}$$

Here, $\hat{\sigma}_z = |e\rangle\langle e| - |g\rangle\langle g|$, $\hat{\sigma}_+ = |e\rangle\langle g|$, and $\hat{\sigma}_- = |g\rangle\langle e|$ are the Pauli operators of the ion's two internal states $|g\rangle$ and $|e\rangle$ (with the transition frequency ω_a between them); Ω_0 is the Rabi frequency describing the strength of the coupling between the applied laser and the ion; η is the Lamb-Dicke (LD) parameter, describing the strength of the coupling between the external and internal states of the ion; and ω_l and ϕ_l are the effective frequency and initial phase of the applied laser fields, respectively. If the LD parameter η is sufficiently small, then the Hamiltonian in Eq. (3) can be approximated to

$$\hat{H}_{\rm L}' = \frac{\hbar\omega_a}{2}\hat{\sigma}_z + \hbar\Omega_0(\hat{\sigma}_+ + \hat{\sigma}_-)\Bigg[(1 + i\eta(\hat{a}_2 + \hat{a}_2^\dagger) \\ -\frac{\eta^2}{2}(\hat{a}_2 + \hat{a}_2^\dagger)^2)e^{-i\omega_l t - i\phi_l} + \text{H.c.}\Bigg] \tag{4}$$

by neglected the higher-order terms $O(\eta^3)$.



FIG. 2: (Color online) Occupancies $|\beta|^2$ in Eq. (8) versus gt for different detunings: $\Delta/g = 0$, $\Delta/g = 2$, $\Delta/g = 5$, and $\Delta/g = 10$. The results indicate that the probability, of the SWAP between the vibrational quanta of the two ions, enhances with the decrease of detuning Δ.

Consequently, the total Hamiltonian of the present two-ion system $\hat{H} = \hat{H}_{\rm ex} + \hat{H}_{\rm L}'$ can be effectively approximated as

$$\hat{H}' = \frac{\hbar\delta_{\rm in}}{2}\hat{\sigma}_z + \hbar\tilde{\Omega}\hat{F} + \hbar(\delta_{\rm ex} - \Omega\hat{F})\hat{a}_2^\dagger\hat{a}_2 - \hbar g(\hat{a}_1\hat{a}_2^\dagger + \hat{a}_1^\dagger\hat{a}_2), \tag{5}$$

in the rotating framework defined by the unity operator $\hat{U}_1 = \exp[-it\omega_l\sigma_z/2 - it\sum_{j=1}^{2}(\nu_1\hat{a}_j^\dagger\hat{a}_j + \nu_j/2)]$. Above, $\hat{F} = \exp(-i\phi_l)\hat{\sigma}_+ + \exp(-i\phi_l)\hat{\sigma}_-$, $\delta_{\rm ex} = \nu_2 - \nu_1$, $\delta_{\rm in} = \omega_a - \omega_l$, $\tilde{\Omega} = 1 - \eta^2/2$, $\Omega = \Omega_0\eta^2$, and $g = 2\mathcal{K}\xi_1\xi_2/d^2$. Suppose further that the ion is driven resonantly, i.e., $\delta_{\rm in} = 0$, then the Hamiltonian in Eq. (5) reduces to

$$\hat{H}'' = \hbar(\delta_{\rm ex} - \Omega\hat{F})\hat{a}_2^\dagger\hat{a}_2 - \hbar g(\hat{a}_1\hat{a}_2^\dagger + \hat{a}_1^\dagger\hat{a}_2) \tag{6}$$

in another rotating framework defined by the unity operator $\hat{U}_2 = \exp(-it\tilde{\Omega}\hat{F})$. Specially, if the effective phase of the laser beams is set as $\phi_l = 0$ and the internal state of the ion is prepared in the eigenstate: $|\varphi\rangle = (|g\rangle + |e\rangle)/\sqrt{2}$ of the operator $\hat{F}$, then the Hamiltonian in Eq. (6) reduces to

$$\hat{H}_{\rm eff} = \hbar\Delta\hat{a}_2^\dagger\hat{a}_2 - \hbar g(\hat{a}_1\hat{a}_2^\dagger + \hat{a}_1^\dagger\hat{a}_2) \tag{7}$$

with $\Delta = \delta_{\rm ex} - \Omega$ being the laser-induced controllable detuning between the vibrations. In the previous works [14–16] switchable couplings between the ions were implemented by adiabatically manipulating the vibrational frequencies ν_j (i.e., $\delta_{\rm ex}$). Here, we fixed ν_j (and thus $\delta_{\rm ex}$ is unchanged) but just adjust the laser-induced effective Rabi frequency Ω to manipulate Δ. Since Ω is relatively-easily controlled by the applied laser and the adiabatic condition is unnecessary to be satisfied. As a consequence, the laser-induced ion-ion coupling/decoupling could be relatively-fast implemented.

The effective Hamiltonian in Eq. (7) directly yields the dynamics for the bosonic modes of the ions

$$\begin{array}{l} |0\rangle_1|1\rangle_2 \longrightarrow \alpha|0\rangle_1|1\rangle_2 + \beta|1\rangle_1|0\rangle_2, \\ |1\rangle_1|0\rangle_2 \longrightarrow \alpha|1\rangle_1|0\rangle_2 + \beta|0\rangle_1|1\rangle_2 \end{array} \tag{8}$$

with

$$\alpha = \left(\frac{1}{2} - \frac{\Delta}{4c}\right)e^{i(c-\Delta/2)t} + \left(\frac{1}{2} + \frac{\Delta}{4c}\right)e^{-i(c+\Delta/2)t}, \tag{9}$$

$$\beta = \frac{g}{c} e^{i(\pi - \Delta t)/2} \sin(ct), \tag{10}$$

and $c = \sqrt{(\Delta/2)^2 + g^2}$. Above, only the two lowest-energy vibrational states of each ion are considered and thus the dynamics is limited within the invariant subspace: $\{|0\rangle_1|1\rangle_2, |1\rangle_1|0\rangle_2\}$, with $|0\rangle_j$ and $|1\rangle_j$ being the ground- and first-excitation states of the jth ion's vibration. The complex parameters α and β satisfy the normalized condition $|\alpha|^2 + |\beta|^2 = 1$, where $|\beta|^2$ is just the probability of the SWAP between the vibrational quanta of the ions. Fig. 2 shows how the occupancy $|\beta|^2$ changes with gt for different Δ, i.e., $\Delta = 0, 2g, 5g$, and $10g$, respectively. When $t\sqrt{(\Delta/2)^2 + g^2} = n\pi/2$ with $n = 0, 1, 2, ...$, the value of $|\beta|^2$ reaches its maximum (amplitude), the maximum probability of the desirable SWAP. It is seen that such an amplitude enhances with the decrease of detuning Δ. Therefore, one can turn on or off the coupling between the ionic vibrations by setting $\Delta \approx 0$ or $\Delta \gg g$, correspondingly. Specifically, such a switchable coupling can be implemented as: Initially, the two ions are trapped in separate wells with the detuning $\Delta = \delta_{\rm ex} \gg g$, i.e., their vibrations are decoupled (shown, e.g., in Fig. 1). Then, a laser pulse with the effective Rabi frequency $\Omega = \delta_{\rm ex}$ is applied to one of the two ions for realizing the resonant coupling (i.e., $\Delta = 0$). The induced evolution reads [16]

$$\hat{U}_{1,2}(gt) = \begin{pmatrix} 0 & \cos(gt) \\ i\sin(gt) & 0 \end{pmatrix}. \tag{11}$$

After this laser driving the vibrations of the ions return to their initial decoupled case.

In the above analysis we have assumed $g/\gamma \gg 1$ and neglected the dissipation (with the rate γ) from the environment. Based on the experimental demonstrations [14, 15], we consider that two ^{40}Ca$^+$ ions are trapped in different potential wells (separated by the distance $d = 40$ μm), and their vibrational frequencies are set as $\nu_1 = 5$ MHz and $\nu_2 = 5.1$ MHz. In this case, $g \approx 11$ KHz and $\delta_{\rm ex} = \nu_2 - \nu_1 = 0.1$ MHz, which means that the two ions are effectively decoupled from each other. Also, the typical decoherent rate of the ionic vibration is $\gamma_{\rm ex} \sim 1$ KHz [14, 15], and the selected electronic levels, $|g\rangle = |{\rm S}_{1/2}\rangle$ and $|e\rangle = |{\rm D}_{5/2}\rangle$, are long-lived with the decoherent rate $\gamma_{\rm in} \approx 1$ Hz [20]. Therefore, the condition $g/\gamma \gg 1$ required in our proposal is satisfied. Since the electric dipole transition between $|{\rm S}_{1/2}\rangle$ and $|{\rm D}_{5/2}\rangle$ is forbidden, here we consider the quadrupole transition (at 729 nm) between them. Consequently, the LD parameter is calculated as $\eta \approx 0.1$, working within the LD regime. Experimentally, a laser beam of power $P = 140$ mW can generate a quadrupole transition between $|{\rm S}_{1/2}\rangle$ and $|{\rm D}_{5/2}\rangle$ with Rabi frequency $\Omega_0 = 1.5$ MHz [21]. This indicates that the desirable effective Rabi frequency $\Omega = \Omega_0\eta^2 = \delta_{\rm ex} \approx 0.1$ MHz [for implementing the operation (11)] could be experimentally realized by applying the laser beam with proper power, as $\Omega_0 \propto \sqrt{P}$.

With the above laser-induced switchable coupling we now discuss how to implement the universal quantum gates by non-adiabatically manipulations. For generality we consider here $n\,(\geq 2)$ potential wells (the adjacent potentials are separated by an uniform distance d), as shown in the Fig. 3. Each potential well confines two ions [15], e.g., Q_n and A_n. The ion Q_n (e.g., a ^{9}Be$^+$ [14]) is used to encode the qubits by their long-lived internal states $|\downarrow\rangle$ and $|\uparrow\rangle$. While, the auxiliary ion A_n (e.g., a ^{40}Ca$^+$ [15]) is used for two purposes: (i) cooling the CM mode $|\psi_{\rm cm}\rangle_n$ of the two ions Q_n and A_n confined in the same trap, and (ii) implementing the switchable coupling between the CM modes of the ions trapped in adjacent potential wells.



FIG. 3: (Color online) Sketch for implementing quantum logic operations with $n\,(\geq 2)$ potential wells. The distance between the adjacent wells is d. Each potential well confines two ions: the red ion (e.g. the ion Q_n) is used to encode the qubit, and the blue ion (e.g., the ion A_n) is used to cool the CM modes of the ions and implement the switchable coupling between the ionic vibrations in the adjacent potential wells.

It has been well-known that the single-qubit operations can be exactly implemented by driving various resonant Raman transitions [22]. The present purpose is to implement the two-qubit controlled-NOT (CNOT) gate [23]

$$\hat{C}_{1,n} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \tag{12}$$

between the qubits (e.g., the ions Q_1 and Q_n) trapped in the different potential wells. For simplicity, we suppose that all the CM modes are initially decoupled and cooled in the vacuum states $\prod_{j=1}^{n} |0_{\rm cm}\rangle_j$. Then, the desirable CNOT gate $\hat{C}_{1n}$ could be realized by the following operational sequence

$$\hat{C}_{1,n} = \hat{V}_1 \hat{M}_{n,1} \hat{S}_n \hat{M}_{1,n} \hat{V}_1. \tag{13}$$

Here, $\hat{V}_1$ represents the operation $|0_{\rm cm}, \uparrow\rangle_1 \longrightarrow \exp[i(1-n)\pi/2]|1_{\rm cm}, \downarrow\rangle_1$ applied to the ion Q_1 for transferring the quantum information (QI) stored in the ion's internal states to the CM mode of the ions trapped in the first well. Experimentally, the operation $\hat{V}_1$ could be realized by applying a red-sideband laser pulse to the ion, see, e.g., Ref. [24]. The operation $\hat{M}_{1,n} = \prod_{j=1}^{n-1} \hat{U}_{(n-j),(n-j+1)}(\pi/2)$ represents $n-1$ resonant pulses applied sequentially on the auxiliary ions A_2, A_3, $\cdots$, and A_n to couple the CM modes in the separated potential wells one by one. Physically, the operation $\hat{M}_{1,n}$ transfers the QI stored in the CM mode of the potential well 1 to that in the distant well n. Similarly, $\hat{M}_{n,1} = \prod_{j=1}^{n-1} \hat{U}_{j,(j+1)}(\pi/2)$ transfers the QI from the potential well n to the first well. Finally, $\hat{S}_n$, a single-ion CNOT gate between the CM mode and internal states of the ion Q_n, can be implemented by applying three sequential laser pulses to the ion (as the demonstrations in the experiment [22]).

One of the advantages of the above non-adiabatic manipulations is that our CNOT gate could be implemented within a significantly-short operational times. First, the durations of operation $\hat{V}_1$ and gate $\hat{S}_n$ by using the Raman schemes of experiment [22] are $t_{\rm v} \approx 8\,\mu$s and $t_{\rm s} \approx 50\,\mu$s, respectively. Second, for the typical parameters [14–16]: $d \approx 40\,\mu$m (the distance between the adjacent potential wells) and $\nu_j \approx 5$ MHz (the vibrational frequencies of the CM modes), the coupling strength between the adjacent CM modes is calculated as $g_{\rm cm} \approx 35$ KHz, and thus the duration of the operation $\hat{U}_{j,(j+1)}(\pi/2)$ is $t_{\rm u} \approx 45\,\mu$s. Obviously, all these durations (i.e., $t_{\rm v}$, $t_{\rm s}$, and $t_{\rm u}$) are significantly shorter than the coherent times (which is about 1 ms) of the CM modes [14, 15]. Note that the laser coolings of CM modes performed by using the auxiliary ions do not affect the qubits. Therefore, during the operation V_1, $\hat{S}_n$, or $\hat{U}_{j,(j+1)}$ applied on the ions 1, n, or j and $j+1$, the other ions can be re-cooled. This implies that the practical limit for the implementation of CNOT gate (12) is the decoherence from the states $|\downarrow\rangle$ and $|\uparrow\rangle$, i.e., the qubit. Fortunately, the coherent times between the selected internal states $|\downarrow\rangle$ and $|\uparrow\rangle$ are relative long, e.g, up to ~ 10 s [22]. This provides us to implement the desirable gates between a large number of trapped ions, e.g., $n \sim 100$. Indeed, the total duration $t_{\text{total}} = 2[t_{\rm v} + (n-1)t_{\rm u}] + t_{\rm s} \approx 9$ ms for generating the CNOT gate (12) is far shorter than the coherent time of the states $|\downarrow\rangle$ and $|\uparrow\rangle$. Based on these estimations, a large number of ions could be entangled by a series of single-qubit operations and CNOT gates, in principle.

In conclusion, we have proposed an improved approach to implement the universal quantum gates between the separately trapped ions. Instead of the previous adiabatic operations of potential wells, here the laser beams are utilized to non-adiabatically manipulate the ion-ion couplings. Consequently, the desirable quantum gates between the ions could be fast implemented by using only a few of laser pulses. A key feature of our proposal is that it could be conveniently generalized to the scaling ion-traps.

Acknowledgements: This work was partly supported by the National Natural Science Foundation of China Grants No. 10874142 and No. 90921010, the Major State Basic Research Development Program of China Grant No. 2010CB923104 (through Program No. 973), and the open project of State Key Laboratory of Functional Materials for Informatics.

---

[1] N. A. Gershenfeld and I. L. Chuang, Science **275**, 350 (1997).

[2] Y. Makhlin, G. Schön, and A. Shnirman, Rev. Mod. Phys. **73**, 357 (2001).

[3] D. Loss and D. P. DiVincenzo, Phys. Rev. A **57**, 120 (1998).

[4] J. M. Raimond, M. Brune, and S. Haroche, Rev. Mod. Phys. **73**, 565 (2001); J. I. Cirac, P. Zoller, H. J. Kimble, and H. Mabuchi, Phys. Rev. Lett. **78**, 3221 (1997).

[5] T. D. Ladd, F. Jelezko, R. Laflamme, Y. Nakamura, C. Monroe, and J. L. O′Brien, Nature (London) **464**, 45 (2010).

[6] D. J. Wineland, C. Monroe, W. M. Itano, D. Leibfried, B. E. King, and D. M. Meekhof, J. Res. Natl. Inst. Stand. Technol. **103**, 259 (1998).

[7] T. Monz, P. Schindler, J. T. Barreiro, M. Chwalla, D. Nigg, W. A. Coish, M. Harlander, W. Hänsel, M. Hennrich, and R. Blatt, Phys. Rev. Lett. **106**, 130506 (2011).

[8] J. I. Cirac and P. Zoller, Phys. Rev. Lett. **74**, 4091 (1995).

[9] K. Mølmer and A. Sørensen, Phys. Rev. Lett. **82** 1835 (1999); G. J. Milburn, S. Schneider, and D. F. V. James, Fortschr. Phys. **48** 801 (2000).

[10] H. Häffner, C. F. Roos, and R. Blatt, Phys. Rep. **469**, 155 (2008).

[11] J. I. Cirac and P. Zoller, Nature (London) **404**, 579 (2000).

[12] D. Kielpinski, C. Monroe, and D. J. Wineland, Nature (London) **417**, 709 (2002).

[13] J. P. Home, D. Hanneke, J. D. Jost, J. M. Amini, D. Leibfried and D. J. Wineland, Science **325**, 1227 (2009).

[14] K. R. Brown, C. Ospelkaus, Y. Colombe, A. C. Wilson, D. Leibfried, and D. J. Wineland, Nature (London) **471**, 196 (2011).

[15] M. Harlander, R. Lechner, M. Brownnutt, R. Blatt, and W. Hänsel, Nature (London) **471**, 200 (2011).

[16] M. Zhang and L. F. Wei, Phys. Rev. A, **83** 064301 (2011).

[17] T. Sleator and H. Weinfurter, Phys. Rev. Lett. **74**, 4087 (1995); D. P. DiVincenzo, Phys. Rev. A **51**, 1015 (1995).

[18] C. A. Blockley, D. F. Walls, and H. Risken, Europhys. Lett. **17**, 509 (1992); D. Leibfried, R. Blatt, C. Monroe, and D. Wineland, Rev. Mod. Phys. **75**, 281 (2003).

[19] L. F. Wei, Y. X. Liu, and F. Nori, Phys. Rev. A, **70** 063801 (2004); L. F. Wei, M. Zhang, H. Y. Jia, and Y. Zhao, Phys. Rev. A, **78** 014306 (2008).

[20] Ch. Roos, Th. Zeiger, H. Rohde, H. C. Näerl, J. Eschner, D. Leibfried, F. Schmidt-Kaler, and R. Blatt, Phys. Rev. Lett. **83**, 4713 (1999).

[21] U. G. Poschinge, G. Huber, F. Ziesel, M. Deiß, M. Hettrich, S. A. Schulz, K. Singer, G. Poulsen, M. Drewsen, R. J. Hendricks, and F. Schmidt-Kaler, J. Phys. B: At. Mol. Opt. Phys. **42**, 154013 (2009).

[22] C. Monroe, D. M. Meekhof, B. E. King, W. M. Itano, and D. J. Wineland, Phys. Rev. Lett. **75**, 4714 (1995).

[23] L. F. Wei, S. Y. Liu, and X. L. Lei, Phys. Rev. A, **65** 062316 (2002).

[24] D. M. Meekhof, C. Monroe, B. E. King, W. M. Itano, and D. J. Wineland, Phys. Rev. Lett. **76**, 1796 (1996).